LOCAL NOXIOUS WEEDS

Spotted Knapweed

(Centaurea maculosa)



Roots: taproot, stout

Stems: erect or ascending, 1 to several, branching above, ridged

Leaves: alternate, surfaces nearly hairless

Flowers: pink to purple disk florets

Origin: Eurasia

Musk Thistle (Carduus nutans)



Roots: taproot, fleshy, stout

Stems: erect, highly branched, with spiny wings

Leaves: alternate, blades simple, margins deeply serrate, surfaces without hair **Flowers:** rose-purple to white disk florets, developing from outer edge to center

Origin: Eurasia

Plumeless Thistle

(Carduus acanthoides)



Roots: taproot, fleshy, stout

Stems: erect, freely branching above, with spiny wings

Leaves: alternate, blades simple, irregular and deeply pinnatifid

Flowers: purple disk florets

Origin: Eurasia

Field Bindweed

(Convolvulus arvensis)



Roots: extensive root system (to a depth of 10m) rhizomes, creeping

Stems: vine, twining, trailing on the surface or climbing **Leaves:** alternate, blades simple, variable, ovate to oblong

Flowers: white or pink corolla, funnel shaped

Origin: Eurasia

Common Mullein

(Verbascum thapsus)



Roots: taproot

Stems: erect, stout, unbranched or a few upright branches near the top **Leaves:** alternate and rosette, basal blades simple, surfaces wooly

Flowers: sulfur-yellow to pale yellow corolla

Origin: Europe