COMPARING COVID-19 VACCINES

COVID-19 vaccines are <u>safe</u>, <u>effective</u>, and free! After you've been fully vaccinated, you can <u>participate in many of the activities</u> that you did prior to the pandemic. Learn about <u>booster shots</u>. Learn about <u>COVID-19 vaccines for children and teens</u>.

<u>and teens</u> .	Pfizer-BioNTech	Moderna	Janssen (<u>view updates</u>)
Vaccine Type	mRNA (FDA Approved)	mRNA	Vector
Age requirement	5 years and older	18 years and older	18 years and older
Number of Doses	Two doses are needed, 21 days apart (or up to six weeks apart, if needed) Some protection provided after the first dose	Two doses are needed, 28 days apart (or up to six weeks apart, if needed) Some protection provided after the first dose	One dose is needed Protection provided two weeks after vaccination
Additional Dose	Recommended for moderately to severely immunocompromised people Given 4 weeks after second shot	Recommended for moderately to severely immunocompromised people Given 4 weeks after second shot	Not recommended at this time
Booster Dose	Recommended for some people who are at higher risk for COVID-19 exposure or severe illness Given 6 or more months after second shot	Recommended for some people who are at higher risk for COVID-19 exposure or severe illness Given 6 or more months after second shot	All those <u>18 years or</u> older are eligible for a booster Given at least 2 months after your first shot
You may choose which COVID-19 vaccine you receive as a booster shot.			
Side Effects	Injection site pain, fatigue, headache, muscle pain, chills, joint pain, fever, nausea, feeling unwell and swollen lymph nodes	Injection site pain, fatigue, headache, muscle pain, chills, joint pain, swollen lymph nodes in the arm that was injected, nausea, vomiting and fever	Injection site pain, headache, fatigue, muscle pain, chills, fever and nausea

If you have any questions or concerns about getting a COVID-19 vaccine, talk to your doctor.



doh.sd.gov/covid/